



**Bart Eastwood D.O.
250 South Main St.
Suite 224A
Blacksburg, VA 24060
540-552-7133**

Post-Op Peroneal tendon repair Protocol

Week 0-2 (Phase 0)

- Weight bearing: No weight bearing. Use crutches/walker/Roll-a-bout as necessary.
- Dressing/wound care: Keep splint on at all times.
- Activity: straight leg raises, toe curls

Week 2-6 (Phase 1: Gentle range-of-motion)

- Weight bearing: May begin very mild partial weight bearing with assistance of crutches/walker only while using CAM boot. You should wear the boot at all times including while sleeping. May advance to weight bearing as tolerated week 4 as long as you can walk without pain. Continue to use at least one crutch, however, for support.
- Wound care: you may get wet very briefly to wash. Keep clean and dry. Report any increasing redness or drainage. You may start gentle massage to skin around the incision starting week 4.
- Activity: Straight leg lift, standing hamstring curl. No ankle motion until week 4, then may begin gentle ankle flexion-extension only, no inversion/eversion.

Week 6-10 (phase 2: initiate strengthening)

Weight bearing: Ween CAM boot, weight bearing as tolerated in lace-up brace.

Wound Care: Continue daily gentle massage to incision site and surrounding skin and tissue. Call if any redness or drainage.

Activity: Full ankle ROM including inversion and eversion. May begin stationary bike and elliptical with the lace up brace.

Week 10-16 (phase 3: advanced strengthening)

Weight bearing: may begin to ween the lace up brace as strength returns.

Wound care: continue daily gentle massage to incision site and surrounding skin and tissue. Call if any increasing redness or drainage.

Activity: begin more advanced strengthening and proprioception. May return to impact activities in week 10.

Week 16 (phase 4: return to sport)

Weight bearing: full weight bearing in regular shoe

Activity: Progressive running. Progressive agility and proprioception. Advancement to plyometrics.